

MONITORING FISH SIZE BY STEREO IMAGING AND PREDICTING GROWTH WITH DES + MACHINE LEARNING

A framework for Aquaculture Digital Twins

From Knowledge to Competitiveness

Portugal-US Partnerships 2026 Summit

Portugal Program



Luiz Claudio Navarro^{1,2}

Lnavarro@ciimar.up.pt

Supervisor(s): Rodrigo Ozorio¹, Aníbal Matos², Anderson Rocha³

1. CIIMAR, University of Porto, Portugal
2. DEEC FEUP, University of Porto, Portugal
3. Recod.ai UNICAMP, Brazil

under the Doctoral Grant 2022 MPP2030-FCT PhD Grant PRT/BD/154260/2022


Data Science

1 Why a tank-specific Digital Twin?

Weight dispersion is challenging in tanks of aquaculture of European seabass (*D. labrax*) and rainbow trout (*O. mykiss*) that are key farmed species in Portugal.


- It drives competition, aggression, poor feed efficiency, and unpredictable harvests.
- Corrective grading is laborious and stressful. Tanks differ in rearing parameters.
- Digital Twins, virtual models synchronized with physical data, predict future states.

This work develops their two kernels: observation and forecasting.



2 Observation kernel

Above-water stereo vision and environmental sensors [1]



Development and production pipelines: stereo images to length and weight distributions [1].

- **Optimized calibration:** validation error < 3.3%; fish length error < 4.2%.
- **Refraction** corrected in two vertical planes; YOLOv11x segments the water line in border.
- **Multi-fish YOLOv11x contours** and a 33-point descriptor drive stereo matching, triangulation, and morphological filtering.
- **Reciprocal KNN stereo matching** through polar descriptor vector.
- **Gradient Boosting** allometry converts length and width to weight.

2.3% | 1.2%
median length MAPE
seabass | trout


6.9% | 4.9%
median weight MAPE
seabass | trout

97.2%
mask mAP50
segmentation

100-day seabass and 71-day trout trials; turbidity reached the 200 NTU sensor limit.

3 Forecasting kernel

Discrete Event System (DES) and Machine Learning [2]



Production cycle, feed-fish-water model, and DES-to-ML methodology [2].

- **DES** simulates individual feed intake, growth, dominance, and metabolism, plus water chemistry.
- **Random Forest surrogates**, trained on 11,029 simulated days, reproduce the DES outputs with $R^2 > 0.996$.
- **Tank-specific hyperparameters** are fitted to the sampled weight distributions.
- **Rapid what-if evaluation** of feeding, grading, and harvesting scenarios.


1.1%
final mean weight
MAPE

8.1%
final weight SD
MAPE

20,000x
RF model faster than
the DES


Validated on 21 European seabass tanks from two feeding trials.

4 Kernels as Digital Twin infrastructure



Each kernel was validated independently; full autonomous integration is the next step.

5 Impact



Conclusion Non-invasive, population-level state estimation linked to tank-specific predictive modeling provides both computational kernels of a Precision Fish Farming Digital Twin, supporting proactive feeding, grading, transfer, and harvest decisions.

[1] L.C. Navarro et al. High-accuracy, non-invasive fish biometrics via above-water stereo vision for biomass and weight dispersion estimation. Computers and Electronics in Agriculture, under review (2026).

[2] L.C. Navarro et al. Predicting weight dispersion in seabass aquaculture using Discrete Event System simulation and Machine Learning modeling. Aquaculture Reports 38 (2024) 102315. <https://doi.org/10.1016/j.aqrep.2024.102315>

Funded by:

